

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
24 June 2004 (24.06.2004)

PCT

(10) International Publication Number
WO 2004/052606 A2

- (51) International Patent Classification⁷: **B29B**
- (21) International Application Number:
PCT/EP2003/013907
- (22) International Filing Date: 8 December 2003 (08.12.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
02447247.4 6 December 2002 (06.12.2002) EP
- (71) Applicants (*for all designated States except US*): **INNO-GENETICS N.V.** [BE/BE]; Technologiepark 6, B-9052 Ghent (BE). **ROCHE DIAGNOSTICS GMBH** [DE/DE]; Sandhofer Strasse 116, 68305 Mannheim (DE).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **CLAEYS, Sofie** [BE/BE]; Briersweg 365, B-8310 Brugge-Sint-Kruis (BE). **JANNES, Geert** [BE/BE]; E. Vanhoorenbekeaan 23/1, B-3010 Leuven (BE). **HABERHAUSEN, Gerd** [DE/DE]; Kapellenwiese 35, 82377 Penzberg (DE). **EMRICH, Thomas** [DE/DE]; Blombergstrasse 7, 82393 Iffeldorf (DE). **VERDOODT, Lia** [BE/BE]; Wielewaalpark 10, B-9840 De Pinte (BE).
- (74) Common Representative: **INNOGENETICS N.V.**; Intellectual Property Department, Industriepark Zwijnaarde 7, Box 4, B-9052 Ghent (BE).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 2004/052606 A2

(54) Title: DETECTION, IDENTIFICATION AND DIFFERENTIATION OF EUBACTERIAL TAXA USING A HYBRIDIZATION ASSAY

(57) Abstract: The present invention relates to a method for the specific detection and/or identification of *Staphylococcus* species, in particular *Staphylococcus aureus*, using new nucleic acid sequences derived from the ITS (Internal Transcribed Spacer) region. The present invention relates also to said new nucleic acid sequences derived from the ITS region, between the 16S and 23S ribosomal ribonucleic acid (rRNA) or rRNA genes, to be used for the specific detection and/or identification of *Staphylococcus* species, in particular of *S. aureus*, in a biological sample. It relates also to nucleic acid primers to be used for the amplification of said spacer region of *Staphylococcus* species in a sample.

DETECTION, IDENTIFICATION AND DIFFERENTIATION OF
EUBACTERIAL TAXA USING A HYBRIDIZATION ASSAY

5 FIELD OF THE INVENTION

The present invention relates to a method for the specific detection and/or identification of *Staphylococcus* species, in particular *Staphylococcus aureus*, using new nucleic acid sequences derived from the ITS (Internal Transcribed Spacer) region.

10 The present invention relates also to said new nucleic acid sequences derived from the ITS region, between the 16S and 23S ribosomal ribonucleic acid (rRNA) or rRNA genes, to be used for the specific detection and/or identification of *Staphylococcus* species, in particular of *S. aureus*, in a biological sample.

It relates also to nucleic acid primers to be used for the amplification of said spacer region of *Staphylococcus* species in a sample.

15

BACKGROUND OF THE INVENTION

20 The genus *Staphylococcus* includes currently 32 described species and 15 subspecies. From the human clinical point of view, *S. aureus* is the most important, but some coagulase-negative species are emerging pathogens especially in nosocomial infections among critical care patients.

Certain species of the *Staphylococcus* genus are more frequently isolated as etiologic agents of a variety of infections in humans. The agents of most concern are *S.*
25 *aureus*, *S. epidermidis*, *S. haemolyticus*, *S. lugdunensis*, *S. warneri* and *S. saprophyticus*.

S. schleiferi has been considered a significant pathogen in some European countries but has only seldom been reported in the US, which demonstrates the variability of the local epidemiology of pathogens.

In veterinary medicine *S. aureus*, *S. intermedius* and *S. hyicus* are the most
30 prominent pathogens

Staphylococcus aureus is one of the most common nosocomial pathogens. It is responsible for several diseases, ranging from superficial cutaneous abscesses to life-threatening intravascular infections. Its propensity to establish prolonged carriage among hospitalized patients and increasing resistance to antibiotics makes control of this

organism within the hospital very difficult.

Knowledge of the epidemiology of *S. aureus* colonization among patients has shed new light on the potential difficulties in interrupting nosocomial transmission. Effective control of *S. aureus* within the hospital and community requires more aggressive measures that include earlier diagnosis of colonized patients, in other words, that include a step of screening.

Because *Staphylococcus* bacteraemia still increases in frequency, it is necessary and urgent to provide more rapid methods of detection and/or identification, using probes and/or primers more sensitive and more specific.

10

SUMMARY OF THE INVENTION

15 It is an object of the present invention to provide new nucleic acid sequences derived from a particular region of the ITS of *Staphylococcus* species, which can be used, for the detection and/or identification of *Staphylococcus* species, in particular of *S. aureus*.

20 The present invention thus provides an isolated nucleic acid molecule consisting of SEQ ID NO 1, the RNA form of said SEQ ID NO 1 wherein T is replaced by U, the complementary form of said SEQ ID NO 1, or any homologue, and the use of said nucleic acid molecule as a target for the detection and/or identification of *Staphylococcus* species.

25

An aspect of the present invention relates to new polynucleotides for use as probes and/or primers, which have as target a particular region of the 16S-23S rRNA spacer region of *Staphylococcus aureus*, and which allow the detection and/or identification of *Staphylococcus* species, in particular of *Staphylococcus aureus*.

30

The present invention thus provides an isolated nucleic acid molecule that specifically hybridizes to SEQ ID NO 1, or to the RNA form of said SEQ ID NO 1 wherein T is replaced by U, or to the complementary form of said SEQ ID NO 1, or to any homologous sequences thereof, or to a fragment of at least 20 contiguous

nucleotides thereof, for the detection and/or identification of *Staphylococcus* species, in particular of *Staphylococcus aureus*.

Another aspect of the present invention relates to sets of probes for the detection
5 and/or identification of *Staphylococcus* species, in particular of *Staphylococcus aureus* in a sample.

Another aspect of the present invention concerns primers allowing specific
amplification of the 16S-23S rRNA spacer region of *Staphylococcus* species, in
10 particular of *S. aureus*.

Another object of the present invention is a composition containing any of the new
sequences of the invention, or any of the new sets of probes and/or primers of the
invention; or a combination thereof.

Another object of the present invention is a kit, in which said probes and/or
15 primers are used, for the detection and/or identification *Staphylococcus* species, in
particular of *Staphylococcus aureus*.

Another object of the present invention is a rapid and reliable hybridization
method for detection and/or identification of *Staphylococcus* species, in particular of
Staphylococcus aureus.

20 Another object of the present invention is a hybridization method based on real
time PCR for detection and/or identification of *Staphylococcus* species, in particular of
Staphylococcus aureus.

TABLE LEGENDS

- 25 Table 1: list of SEQ IDs
Table 2: primer pairs
Table 3: set of probes
Table 4: *Staphylococcus* species

30 DETAILED DESCRIPTION OF THE INVENTION

The following definitions serve to illustrate the terms and expressions used in the
different embodiments of the present invention as set out below.

The terms "spacer" and "TTS" (Internal Transcribed Spacer) are abbreviated terms

both referring to the region between the 16S and 23S rRNA or between the 16S and 23S rRNA genes.

The term "probe" refers to single stranded oligonucleotides or polynucleotides which have a sequence which is sufficiently complementary to hybridize to the target
5 sequence to be detected.

Preferably the probes of the invention are 70%, 80%, 90%, or more than 95% homologous to the exact complement of the target sequence to be detected. These target sequences are either genomic DNA or precursor RNA, or amplified versions thereof.

10 The probes of the invention can be formed by cloning of recombinant plasmids containing inserts including the corresponding nucleotide sequences, if need be by cleaving the latter out from the cloned plasmids upon using the adequate nucleases and recovering them, e.g. by fractionation according to molecular weight.

The probes according to the present invention can also be synthesized chemically,
15 for instance by the conventional phospho-triester method.

The term "complementary" nucleic acids as used herein means that the nucleic acid sequences can form a perfect base-paired double helix with each other.

The terms "polynucleic acid", "nucleic acid", and "polynucleotide" correspond to
20 either double-stranded or single-stranded cDNA or genomic DNA or RNA, containing at least 5, 10, 20, 30, 40 or 50 contiguous nucleotides. A polynucleic acid which is smaller than 100 nucleotides in length is referred to as an "oligonucleotide".

They can also refer to modified nucleotides such as inosine or nucleotides containing modified groups which do not essentially alter their hybridization
25 characteristics.

Single stranded polynucleic acid sequences are always represented in the current invention from the 5' end to the 3' end.

They can be used as such, or in their complementary form, or in their RNA form wherein T is replaced by U.

30 The term "closest neighbor" means the taxon which is known or expected to be most closely related in terms of DNA homology and which has to be differentiated from the organism of interest.

The expression "taxon-specific hybridization" or "taxon-specific probe" means that the probe only hybridizes to the DNA or RNA from the taxon for which it was designed and not to DNA or RNA from other taxa.

5 The term taxon can refer to a complete genus or a sub-group within a genus, a species or even subtype within a species (subspecies, serovars, sequevars, biovars...).

The term "specific amplification" or "specific primers" refers to the fact that said primers only amplify the spacer region from these organisms for which they were designed, and not from other organisms.

10 The term "sensitivity" refers to the number of false negatives: i.e. if 1 of the 100 strains to be detected is missed out, the test shows a sensitivity of $(100-1/100)\% = 99\%$.

The term "specificity" refers to the number of false positives: i.e. if on 100 strains detected, 2 seem to belong to organisms for which the test is not designed, the specificity of the test is $(100-2/100)\% = 98\%$.

15 The oligonucleotides or polynucleotides selected as being "preferential" show a sensitivity and specificity of more than 80%, preferably more than 90% and most preferably more than 95%.

20 The term "solid support" can refer to any substrate to which a polynucleotide probe can be coupled, provided that it retains its hybridization characteristics and provided that the background level of hybridization remains low. Usually the solid substrate will be a microtiter plate, a membrane (e.g. nylon or nitrocellulose) or a microsphere (bead). Prior to application to the membrane or fixation it may be convenient to modify the nucleic acid probe in order to facilitate fixation or improve the hybridization efficiency. Such modifications may encompass homopolymer tailing,
25 coupling with different reactive groups such as aliphatic groups, NH_2 groups, SH groups, carboxylic groups, or coupling with biotin, haptens or proteins.

30 The term "labeled" refers to the use of labeled nucleic acids. Labeling may be carried out by the use of labeled nucleotides incorporated during the polymerization step of the amplification such as illustrated by Saiki et al. (1988) or Bej et al. (1990) or by the use of labeled primers, or by any other method known to the person skilled in the art. The nature of the label may be isotopic (^{32}P , ^{35}S , etc.) or non-isotopic (biotin, digoxigenin, fluorescent dye, biotin, enzyme, etc.).

The term "signal" refers to a series of electromagnetic waves (for example fluorescence), or changes in electrical current which carry information. The signal can be directly visible, or can be made visible and/or interpretable by different means or devices.

5 The "sample" may be any biological material. This biological material may be taken either directly from the infected human being, or animal, or after culturing or enrichment, or from food, from the environment, etc .

Biological material may be for example expectoration of any kind, broncheolavages, blood, skin tissue, biopsies, lymphocyte blood culture material,
10 colonies, etc. Said samples may be prepared or extracted according to any of the techniques known in the art.

The *Staphylococcus* species that are clinically relevant in the meaning of the present invention are *S. aureus*, *S. auricularis*, *S. capitis*, *S. caprae*, *S. cohnii*, *S. epidermidis*, *S. haemolyticus*, *S. hominis*, *S. lugdunensis*, *S. pasteurii*, *S. saccharolyticus*,
15 *S. saprophyticus*, *S. schleiferi*, *S. simulans*, *S. warneri*, and *S. xyloso* (Table 4).

The ITS is already known for some *Staphylococcus* species (WO96/00298).

In further studies, the full genome sequencing of different *Staphylococcus* species
20 has revealed that these organisms contain at least 5 ribosomal RNA operons in their genome.

In particular, within *Staphylococcus* species, *S. aureus* strains show a variety of spacer sequences even within one single isolate.

Those different ITS, more than 16 types of sequences, vary also in length in the
25 range of 300 to 550 base pairs.

To solve the problems generated by this very high variability, the present invention provides a particular region of the ITS, identified and delimited for its great advantage of offering a unique target sequence for the detection and/or identification of all *Staphylococcus* species, and in particular of all *Staphylococcus* species clinically
30 relevant, and more particularly of *S. aureus*.

Indeed, it has been discovered that the target sequence of the invention are found in all type of spacer of every *Staphylococcus* species, in particular of every *Staphylococcus* species that are clinically relevant.

This particular region of the ITS, also referred to as the "target region" or "target sequence", can be defined as a nucleic acid molecule consisting of SEQ ID NO 1 or SEQ ID NO 2, or as a nucleic acid molecule that is homologous to SEQ ID NO 1 or 2, their RNA form wherein T is replaced by U, or their complementary form.

5 This term "target sequence" covers all the homologous sequences found in the ITS of any *Staphylococcus* species, said homologous sequences are also referred to herein after as "homologues". The degree of homology is then higher than 75%, generally higher than 80%, and even higher than 90%.

In the framework of this invention, "homologues" are then homologous sequences
10 to SEQ ID NO 1 or 2 or to any fragment thereof, localized in the ITS region of any *Staphylococcus* species, SEQ ID NO 1 and 2 being derived from different *S. aureus* strains.

New polynucleotides for use as probes and/or primers designed from the target
15 sequence of the invention for the detection and/or identification of *Staphylococcus* species are also an object of the invention.

In other words, an object of the invention relates to new polynucleotides for use as probes and/or primers, which hybridize with the target sequence of the invention for the
20 detection and/or identification of *Staphylococcus* species.

In particular, an object of the invention is an isolated nucleic acid molecule that specifically hybridizes to SEQ ID NO 1 or 2, or to the RNA form of said SEQ ID NO 1 or 2 wherein T is replaced by U, or to the complementary form of said SEQ ID NO 1 or 2, or to a fragment of at least 20 contiguous nucleotides thereof, or to any of their
25 homologues, for the detection and/or identification of *Staphylococcus* species, in particular of *S. aureus*.

Preferred polynucleotide probes are between about 5 to about 50 bases in length, more preferably from about 10 to about 25 nucleotides and are sufficiently homologous to the target sequence.

30 Polynucleotides of SEQ IDs NO 1 to 70 and any of their homologues may be used as probes.

Preferred probes are polynucleotides of SEQ IDs NO 14, 16 to 23, 25 to 32, 35 to 42 and homologues.

Preferred primers of the invention are single stranded DNA polynucleotides capable of acting as a point of initiation for synthesis of the target sequence of the invention. The length and the sequence of a primer of the invention must be such that they allow to prime the synthesis of the extension products.

5 Preferably a primer of the invention is about 5 to about 50 nucleotides long, preferably about 15 to about 25. Its specific length and sequence is to be chosen depending on the conditions used such as temperature and ionic strength.

Preferred primers of the invention amplify the target sequence. In other words, preferred primers of the invention amplify SEQ ID NO 1 or SEQ ID NO 2 and/or
10 homologues.

Preferred primers of the invention are polynucleotides of SEQ IDs NO 51, 52, 53, 55, 58 65, 67, 68, 69, 70, and homologues.

The fact that amplification primers do not have to match exactly with the corresponding template sequence to warrant proper amplification is amply documented
15 in the literature (Kwok et al., 1990).

The amplification method used can be either polymerase chain reaction (PCR; Saiki et al., 1988), ligase chain reaction (LCR; Landgren et al., 1988; Wu & Wallace, 1989; Barany, 1991), nucleic acid sequence-based amplification (NASBA; Guatelli et al., 1990; Compton, 1991), transcription-based amplification system (TAS; Kwok et al.,
20 1989), strand displacement amplification (SDA; Duck, 1990; Walker et al., 1992) or amplification by means of Q β replicase (Lizardi et al., 1988; Lomeli et al., 1989) or any other suitable method to amplify nucleic acid molecules known in the art.

The preferred polynucleotides of the invention for use as primers or as probes are
25 listed in Table 1.

Polynucleotides of the invention may differ in sequence from any of the polynucleotides specified in Table 1, or from any of their homologues, either by addition to or removal from any of their respective extremities of one or several nucleotides, or by changing one or more nucleotides within said sequences, or a combination of both,
30 provided that the equivalents then obtained still hybridize with the target sequence as the corresponding unmodified polynucleotides. Said equivalent polynucleotides share at least 75% homology, preferably more than 80%, most preferably more than 85% homology with the corresponding unmodified polynucleotides.

When using an equivalent of a polynucleotide, it may be necessary to modify the hybridization conditions to obtain the same specificity as the corresponding unmodified polynucleotide.

As a consequence, it will also be necessary to modify accordingly the sequence of
5 other polynucleotides when the polynucleotides are to be used in a set under the same hybridization conditions. These modifications can be done according to principles known in the art, e.g. such as those described in Hames B and Higgins S (Eds): Nucleic acid hybridization. Practical approach. IRL Press, Oxford, UK, 1985.

10 The polynucleotides primers and/or probes of the invention may also comprise nucleotide analogues such as phosphorothioates (Matsukura et al., 1987), alkylphosphorothioates (Miller et al., 1979) or peptide nucleic acids (Nielsen et al., 1991; Nielsen et al., 1993) or may contain intercalating agents (Asseline et al., 1984), etc.

The modified primers or probes require adaptations with respect to the conditions
15 under which they are used in order to obtain the required specificity and sensitivity. However the results of hybridization should remain essentially the same as those obtained with the unmodified polynucleotides.

The introduction of these modifications may be advantageous in order to influence some characteristics such as hybridization kinetics, reversibility of the hybrid-formation,
20 biological stability of the polynucleotide molecules, etc.

The probes and primers of the invention are used in methods, also objects of the present invention, for the detection and/or identification of *Staphylococcus* species, in particular of *S. aureus*.

25 Detection and/or identification of the target sequence can be performed by using a electrophoresis method, a hybridization method or a sequencing method.

A method of the invention for the detection of one or more *Staphylococcus* species in a sample comprises the following steps:

30 - First, and if necessary, the nucleic acids present in the sample are made available for amplification and/or hybridization.

- Secondly, and also if necessary, the nucleic acids, if present, are amplified with one or another target amplification system, as specified below. Usually, amplification is needed to enhance the subsequent hybridization signal. However for some samples, or

for some highly sensitive signal-amplification systems, amplification might not be necessary.

- Thirdly, the nucleic acids present in the sample or the resulting amplified product are contacted with probes, and hybridization is allowed to proceed.

- 5 - Finally, the hybrids are detected using a convenient and compatible detection system. From the hybridization signals or patterns observed the presence or absence of one or several *Staphylococcus* species can be deduced.

The amplification system used may be more or less universal, depending on the specific application needed.

- 10 By using universal primers located in the conserved flanking regions (16S and 23S gene) of the rRNA spacer, the spacer region of most if not all organisms of eubacterial origin will be amplified.

- For some applications it may be appropriate to amplify not all organisms present in the sample but one or several *Staphylococcus* species. This may be achieved using
15 specific primers located in the target region of *Staphylococcus* species, and for example the polynucleotides of SEQ IDs NO 69 and 70 or their homologues may be used as such a primer pair, or preferably the polynucleotides of SEQ IDs NO 58 and 68 or their homologues.

- 20 In particular, a method of the invention for detection and/or identification of *Staphylococcus* species, notably of *Staphylococcus aureus*, in a sample comprises the steps of:

- (i) if need be releasing, isolating and/or concentrating the polynucleic acids in the sample;
- 25 (ii) if need be amplifying the 16S-23S rRNA spacer region, or a fragment comprising the target sequence, or the target sequence or a fragment thereof, with at least one suitable primer pair;
- (iii) hybridizing the polynucleic acids of step (i) or (ii) with at least one polynucleotide probe that hybridizes to the target sequence, wherein the target sequence
30 consists of SEQ ID NO 1 or 2 or homologues thereof, or to their RNA form wherein T is replaced by U, or to their complementary form, or to a fragment of at least 20 contiguous nucleotides thereof,
- (iv) detecting the hybrids formed, and

(v) interpreting the signal(s) obtained and inferring the presence of *Staphylococcus* species and/or identifying the *Staphylococcus* species in the sample.

Preferably, the probes of the inventions hybridize under conditions of high stringency.

Under high stringency conditions only complementary nucleic acid hybrids are formed. Accordingly, the stringency of the assay conditions determines the amount of complementarity needed between two nucleic acid strands forming a hybrid. Stringency is chosen to maximize the difference in stability between the hybrid formed with the target and the non-target nucleic acid.

The hybridization conditions are chosen in such a way that the signal of hybridization obtained when a polynucleotide of the invention hybridizes specifically to a target sequence, is different from the signal obtained when said polynucleotide hybridizes to a target sequence in a non-specific manner.

In practice, the different signals may be visualized for example when its intensity is two, five, ten or more times stronger with a specific hybridization to the target, as compared to non-specific hybridization to the target sequence, LiPA system for example.

The different signals may also be visualized when different peaks are drawn in a melting curve analysis, for instance when using a real time PCR method.

The fragment mentioned in the amplification or the hybridization step of any method of the invention may comprise 20 to 50, 20 to 80 or 20 to 100 contiguous nucleotides of SEQ ID NO 1 or 2 or of any homologues.

In one embodiment, a very convenient and advantageous technique for the detection of target sequences that are possibly present in the sample is the real time PCR method.

There are different formats for the detection of amplified DNA, notably TaqMan™ probes, Molecular Beacons probes, or FRET hybridization probes.

Concerning the TaqMan™ probes, a single-stranded hybridization probe is labeled with two components. When the first component, the so-called fluoescer, is excited with light of a suitable wavelength, the absorbed energy is transferred to the second component, the so-called quencher, according to the principle of fluorescence resonance energy transfer. During the annealing step of the PCR reaction, the hybridization probe binds to the target DNA and is degraded by the 5'-3' exonuclease activity of the polymerase, for example Taq Polymerase, during the elongation phase. As a result the excited fluorescent component and the quencher are spatially separated from one another and thus a fluorescence emission of the first component can be measured (EP B 0 543 942 and US 5,210,015).

Concerning Molecular Beacons probes, the probes are also labeled with a first component and with a quencher, the labels preferably being located at different ends of an at least partially self-complementary probe. As a result of the secondary structure of the probe, both components are in spatial vicinity in solution. After hybridization to the target nucleic acids both components are separated from one another such that after excitation with light of a suitable wavelength the fluorescence emission of the first component can be measured (US 5,118,801).

The Fluorescence Resonance Energy Transfer (FRET) hybridization probe test format is especially useful for all kinds of homogenous hybridization assays (Matthews, J. A. and Kricka, L. J., Anal Biochem 169 (1988) 1-25). It is characterized by two single-stranded hybridization probes which are used simultaneously and are complementary to adjacent sites of the same strand of an (amplified) target nucleic acid. Both probes are labeled with different fluorescent components. When excited with light of a suitable wavelength, a first component transfers the absorbed energy to the second component according to the principle of fluorescence resonance energy transfer such that a fluorescence emission of the second component can be measured only when both hybridization probes bind to adjacent positions of the target molecule to be detected.

When annealed to the target sequence, the hybridization probes must be located very close to each other, in a head to tail arrangement. Usually, the gap between the labeled 3' end of the first probe and the labeled 5' end or the second probe is as small as possible, and notably consists of about 0 to 25 bases, and preferably of about 1 to about 5

bases. This allows for a close vicinity of the FRET donor compound and the FRET acceptor compound, which is typically 10-100 Ångstrom.

Alternatively to monitoring the increase in fluorescence of the FRET acceptor component, it is also possible to monitor fluorescence decrease of the FRET donor
5 component as a quantitative measurement of hybridization event.

Among all detection formats known in the art of real time PCR, the FRET-hybridization probe format has been proven to be highly sensitive, exact and reliable (WO 97/46707; WO 97/46712; WO 97/46714). Yet, the design of appropriate FRET hybridization probe sequences may sometimes be limited by the special characteristics
10 of the target nucleic acid sequence to be detected.

As an alternative to the usage of two FRET hybridization probes, it is also possible to use a fluorescent-labeled primer and only one labeled polynucleotide probe (Bernard, P. S., et al., Anal. Biochem. 255 (1998) 101-7). In this regard, it may be chosen arbitrarily, whether the primer is labeled with the FRET donor or the FRET acceptor
15 compound.

FRET hybridization probes (also called Hybprobes or FRET-probes) can also be used for melting curve analysis (WO 97/46707; WO 97/46712; WO 97/46714). In such an assay, the target nucleic acid is amplified first in a typical PCR reaction with suitable amplification primers. The hybridization probes may already be present during the
20 amplification reaction or be added subsequently. After completion of the PCR-reaction, the temperature of the sample is consecutively increased. Fluorescence is detected as long as the hybridization probe is bound to the target DNA. At the melting temperature, the hybridization probe is released from their target, and the fluorescent signal is decreasing immediately down to the background level. This decrease is monitored with
25 an appropriate fluorescence versus temperature-time plot such that the negative of a first derivative function can be calculated. The temperature value corresponding to the obtained maximum of such a function is then taken as the determined melting temperature of said pair of FRET hybridization probes.

Point mutations or polymorphisms within the target nucleic acid result in a less
30 then 100% complementarity between the target nucleic acid and the FRET probes, thus resulting in a decreased melting temperature. This enables for a common detection of a pool of sequence variants by means of FRET-Hybprobe hybridization, whereas subsequently, different members of said pool may become discriminated by means of performing melting curve analysis.

Instead of FRET hybridization probes, Molecular Beacons may alternatively be used for melting curve analysis.

Upon the availability of Real-Time PCR and homogenous Real-Time PCR melting curve analysis, discrimination of certain types of species or strains became possible using either double stranded DNA binding dyes such as SybrGreenTM, or, alternatively, specifically designed hybridization probes hybridizing to different but similar target sequences.

In the first case, melting temperature of the generated double stranded PCR product has to be determined. Yet, this method has only limited applications since few differences cannot be monitored efficiently, because minor sequence variations only result in subtle melting temperature differences.

Alternatively, hybridization probes may be used in such a way that the melting temperature of the probe/target nucleic acid hybrid is being determined.

There are different real time PCR platforms such as the ABI/PrismTM equipments, and in particular the LightCyclerTM apparatus, all based on the same principle consisting of measuring the light emission, continually monitoring the emission peak during the melt cycle, determining and visualizing the temperatures (melting peaks) at which the labeled probes detach from the amplification products. The melting peak data are characteristic of a particular [probe:target] sequence because mismatches between probe and target affect the kinetics of melting, producing different melting peaks for each species of interest.

The LightCyclerTM platform offers many advantages and in particular a gain of time and the possible use of several different sequence-specific fluorescent probe detection systems such as hybridization probes (HybProbes), TaqManTM probes, Molecular Beacons and biprobes (SYBR Green I).

In a preferred method of the present invention, the HybProbe system is used, consisting of two adjacent polynucleotide probes derived from the target region of the invention, in a head-to-tail orientation, spaced by a few nucleotides, generally 0 to 25, preferably about 1 to about 5. One of the probes is labeled at its 3' end by a donor dye, the other is labeled with an acceptor molecule at its 5' end, and is phosphate blocked at the 3' end (to prevent its acting as a primer). The donor dye is generally fluorescein, and the acceptor molecule generally LC Red640 or 705.

The detection of the target sequence of the invention may be achieved also by an internal labeled PCR strand and a detection probe located on the opposite strand. The signal is dependent on the spatial approximation of the dyes, and is dependent on the amount of the target.

5 When both probes are hybridized to their target sequence the emitted light of the donor is transmitted to the acceptor fluorophore by Fluorescence Resonance Energy Transfer (FRET), and the emitted fluorescence (640 or 705 nm) can be detected. The intensity of the emitted fluorescence increases in parallel with the target DNA, product of the amplification.

10 The LightCycler probes offer the advantage over the TaqManTM probes of not requiring hydrolysis and, therefore, no additional extension of the PCR times (annealing-elongation ≤ 12 s). It is therefore possible to take advantage of the high-speed thermal cycling of the LightCycler, and complete the PCR program in only 45 minutes.

And the most recent generations of this real-time PCR platform are able to monitor
15 several probes in a single reaction, allowing the detection and/or identification of different *Staphylococci*, at the species level and also at lower taxonomical levels.

Moreover, it has been shown that the methods designed for TaqMan technology can be easily converted to HybProbe technology with equivalent results (Haematologica vol. 85 (12) pp. 1248-1254, December 2000).

20 Therefore another object of the invention relates to sets of 2 polynucleotide probes, also referred to as HybProbes, both HybProbes hybridizing to the same target sequence, adjacent to each other, with no more than 25 nucleotides between said 2 HybProbes, preferably with no more than 10 nucleotides, in particular with no more than 5 nucleotides.

25 One of the HybProbes is labeled with an acceptor fluorophore and the other with a donor fluorophore of a fluorescence energy transfer pair such that upon hybridization of the two HybProbes with the target sequence, the donor and acceptor fluorophores are within 0 to 25 nucleotides of one another, and preferably within 0 to 5 nucleotides of one another.

30

For detecting and/or identifying *Staphylococcus* species, in particular *Staphylococcus* species clinically relevant, a set of two polynucleotide probes may be used, said two probes hybridizing to SEQ ID NO 1 or SEQ ID NO 2, or to the RNA form of said SEQ ID NO 1 or 2 wherein T is replaced by U, or to the complementary

form of said SEQ ID NO 1 or 2, or to homologues, wherein there are no more than 25 nucleotides, preferably no more than 5 nucleotides, between said two probes.

A set of probes of the invention may also consist of 3, 4, 5, 6, 7, 8, 9, 10, or more, probes, but it preferably consists of 2 to 5 probes, and more preferably of 2 or 3 probes.

The sets of probes listed in Table 3 and their homologues are preferred sets of the invention.

Sets of 2 polynucleotides, one for use as primer, the other for use as probe, may also be used, both said primer and probe hybridizing to the target sequence consisting of SEQ ID NO 1 or 2, of the RNA form of said SEQ ID NO 1 or 2 wherein T is replaced by U, of the complementary form of said SEQ ID NO 1 or 2, or of any homologues, wherein there are no more than 25 nucleotides, preferably no more than 5 nucleotides, between said primer and said probe.

The sets of at least 2 polynucleotides of the invention are used in methods for the detection and/or identification of *Staphylococcus* species, in particular of *S. aureus*.

A method of the present invention for detection and/or identification of *Staphylococcus* species in a sample, comprises the steps of:

- (i) if need be releasing, isolating and/or concentrating the polynucleic acids in the sample;
- (ii) amplifying the 16S-23S rRNA spacer region, or the target sequence, or a part of the spacer comprising the target sequence, or a part of the target sequence, with at least one suitable primer pair;
- (iii) hybridizing the polynucleic acids with at least one set of at least two HybProbes that hybridize to the target sequence, wherein the target sequence consist of SEQ ID NO 1 or 2, or of the RNA form of said SEQ ID NO wherein T is replaced by U, or of the complementary form of said SEQ ID NO, or of any homologues, or of a fragment of at least 20 contiguous nucleotides thereof;
- (iv) detecting the hybrids formed in step (iii);
- (v) inferring the presence of *Staphylococcus* species, or identifying the *Staphylococcus* species in the sample from the differential hybridization signals obtained in step (iv).

For example, a primer pair used in the amplification step is any combination of a forward primer consisting of SEQ ID NO 45, 49, 50, 52, 56, 61, 63, 64, 65, 66, 67, 68 or their homologues, and a reverse primer consisting of SEQ ID NO 46, 47, 48, 51, 53, 54, 55, 57, 58, 59, 60, 62, or their homologues.

5

For example, a set of 2 HybProbes used in the hybridization step is any combination of 2 HybProbes chosen among polynucleotides of SEQ IDs NO 3 to 70 or their homologues, provided that the gap between said two HybProbes when hybridized to the target sequence is less than 25 nucleotides, preferably less than 5 nucleotides.

10

One of the advantages of the HybProbes system resides in the fact that it allows the detection of sequence variation, including mutations, polymorphisms and other variant nucleic acid species, based on the following molecular concept: one of the HybProbe is a tightly binding "anchor probe" whereas the adjacent "sensor probe" spans the region of sequence variation. During melting of the final PCR product, the sequence alteration is detected as a change in the melting temperature (T_m) of the sensor probe.

15

For example, if the sample contains only SEQ ID NO 1, using Hybprobes that specifically hybridize to said SEQ ID NO 1 would generate a single melting peak. If there is also a homologue in the sample, using the same two HybProbes would generate two peaks, as far as there is one mismatched base which generally induces a temperature shift easily observable.

20

Depending on the polynucleotides selected, their T_m and the hybridization conditions, the fluorescence may be measured during the amplification step, generating then amplification curves, or after the amplification step, for a melting curve analysis generating melting curves.

25

Thus the signal obtained may be visualized in the form of amplification curves or in the form of melting curves, from which it is possible to infer the presence of *Staphylococcus* species, and/or to infer which one(s) of the *Staphylococci* are present.

30

In particular, a method for detection and/or identification of *Staphylococcus* species in a sample comprises also the steps of

(i) if need be releasing, isolating and/or concentrating the polynucleic acids in the sample, and

- (ii) amplifying the target sequence, or a part of it, with a primer pair that is labeled,
- (iii) hybridizing the polynucleic acids with at least one HybProbe that hybridize, adjacent to said labeled primer with less than 25 nucleotides in between, to SEQ ID NO 1, or to the RNA form of said SEQ ID NO 1 wherein T is replaced by U, or to the complementary form of said SEQ ID NO 1, or to any homologues, or to a fragment of at least 20 contiguous nucleotides thereof,
- (iv) detecting the hybrids formed, and
- (v) inferring the presence of *Staphylococcus* species, and/or identifying the *Staphylococcus* species in the sample from the signals obtained in step (iv).

A method of the invention using the HybProbes system, may be adapted for the detection and identification of *Staphylococcus aureus*, allowing the distinction of *S. aureus* from other species, and in particular from coagulase negative *Staphylococci* (CoNS).

Then, in the amplification step, suitable primers are primer pairs that specifically amplify the target sequence which consists of SEQ ID NO 1, or of the RNA form of said SEQ ID NO wherein T is replaced by U, or of the complementary form of said SEQ ID NO.

In the hybridization step, the HybProbes should hybridize specifically to SEQ ID NO 1 or 2, or to the RNA form wherein T is replaced by U, or to the complementary form.

Therefore, *S. aureus* strains can be unequivocally distinguished from all other organisms examined by melting curve analysis.

Furthermore, only CoNS give rise to melting peaks; no relevant signals are obtained with non-*Staphylococci* or human genomic DNA.

Preferred primer pairs used in this particular example are any combinations of forward primers chosen among SEQ ID NO 68 or 69 or their homologues and reverse primers chosen among SEQ ID NO 58 or 70 or their homologues.

The sets of HybProbes listed in Table 3 or their homologues are the preferred sets of HybProbes of the invention. A more preferred set of 2 Hybprobes consists of SEQ ID NO 17 or homologues and SEQ ID NO 19 or homologues.

The set of HybProbes consisting of SEQ ID NO 17 and 19 is able to detect *S. aureus*, *S. epidermidis*, and *S. haemolyticus* with a high sensitivity.

Each polynucleotide listed in Table 1, corresponding to SEQ ID NO 1 to SEQ ID NO 70 and any of their homologues, may be used in any methods of the present invention as a primer and/or as a probe, alone or in combination.

5

A second embodiment based also on a hybridization method is the Line Probe Assay technique. The Line Probe Assay (LiPA) is a reverse hybridization format (Saiki et al., 1989) using membrane strips onto which several polynucleotide probes (including
10 negative or positive control polynucleotides) can be conveniently applied as parallel lines.

The LiPA technique, as described by Stuyver et al. (1993) and in international application WO 94/12670, provides a rapid and user-friendly hybridization test. Results can be read within 4 h. after the start of the amplification. After amplification during
15 which usually a non-isotopic label is incorporated in the amplified product, and alkaline denaturation, the amplified product is contacted with the probes on the membrane and the hybridization is carried out for about 1 to 1,5 h. Consequently, the hybrids formed are detected by an enzymatic procedure resulting in a visual purple-brown precipitate. The LiPA format is completely compatible with commercially available scanning devices,
20 thus rendering automatic interpretation of the results possible. All those advantages make the LiPA format liable for use in a routine setting.

The LiPA format is an advantageous tool for detection and/or identification of pathogens at the species level but also at higher or lower taxonomical levels. For instance, probe-configurations on LiPA strips can be selected in such a manner that they
25 can detect the complete genus of *Staphylococcus* or can identify species within the genus (e.g. *Staphylococcus aureus*, *epidermidis*, etc) or can in some cases even detect subtypes (subspecies, serovars, sequevars, biovars, etc. whatever is clinically relevant) within a species.

The ability to simultaneously generate hybridization results with a large number of
30 probes is another benefit of the LiPA technology. In many cases the amount of information which can be obtained by a particular combination of probes greatly outnumbers the data obtained by using single probe assays. Therefore the selection of probes on the membrane strip is of utmost importance since an optimized set of probes will generate the maximum of information possible.

These probes can be applied to membrane strips at different locations and the result is interpreted as positive if at least one of these probes is positive. Alternatively these probes can be applied as a mixture at the same location, hereby reducing the number of lines on a strip. This reduction may be convenient in order to make the strip more concise or to be able to extend the total number of probes on one strip.

Another alternative approach, in view of its practical benefits, is the synthesis of polynucleotides harboring the sequences of two or more different probes, referred to as degenerate probes, which then can be further processed and applied to the strip as one polynucleotide molecule. This approach would considerably simplify the manufacturing procedures of the LiPA-strips. For example, probes with nucleotide sequences A and B are both required to detect all strains of taxon X. In the latter alternative a probe can be synthesized having the nucleotide sequence AB. This probe will have the combined characteristics of probes A and B.

By virtue of the above-mentioned properties the LiPA system can be considered as an efficient format for a hybridization method wherein several organisms need to be detected simultaneously in a sample.

However, it should be clear that any other hybridization assay, whereby different probes are used under the same hybridization and wash conditions can be used for the above-mentioned detection and/or selection methods. For example, it may be possible to immobilize the target nucleic acid to a solid support, and use mixtures of different probes, all differently labeled, resulting in a different detection signal for each of the probes hybridized to the target. And nowadays many different supports are available.

As an example, the procedure to be followed for the detection of one or more *Staphylococcus* species in a sample using the LiPA format is outlined below:

- First, and if necessary, the nucleic acids present in the sample are made available for amplification and/or hybridization.

- Secondly, the nucleic acids, if present, are amplified with one or another target amplification system, as specified below. Usually, amplification is needed to enhance the subsequent hybridization signal.

- Thirdly, eventually after a denaturation step, the nucleic acids present in the sample or the resulting amplified product are contacted with LiPA strips onto which one or more probes (for example DNA-, RNA-, degenerate or modified probes), allowing the detection of the organisms of interest, are immobilized, and hybridization is allowed to proceed.

- Finally, eventually after having performed a wash step, the hybrids are detected using a convenient and compatible detection system. From the hybridization signals or patterns observed the presence or absence of one or several organisms screened for in that particular biological sample can be deduced.

- 5 The amplification system used may be more or less universal, depending on the specific application needed.

By using universal primers located in the conserved flanking regions of the rRNA spacer, i.e. in the 16S gene and the 23S gene, the spacer region of most if not all organisms of eubacterial origin will be amplified.

- 10 For some applications it may be appropriate to amplify not all organisms present in the sample but more specifically *Staphylococcus* species. This may be achieved using specific primers located in the target region of *Staphylococcus* species, for example the polynucleotides of SEQ IDs NO 69 and 70 or their homologues may be used as such a primer pair.

15

A method of the invention for detection and/or identification of *Staphylococcus* species in a sample, comprises the steps of:

- (i) if need be releasing, isolating and/or concentrating the polynucleic acids present in the sample;
- 20 (ii) if need be amplifying the 16S-23S rRNA spacer region, or a part of it, with at least one suitable primer pair;
- (iii) hybridizing the polynucleic acids with at least one probe that hybridizes to the target sequence consisting of SEQ ID NO 1 or 2, or of the RNA form of said SEQ ID NO 1 or 2 wherein T is replaced by U, or of the complementary form of said
- 25 SEQ ID NO, or of any homologues, or of a fragment of at least 20 contiguous nucleotides thereof;
- (iv) detecting the hybrids formed in step (iii);
- (v) identification of the micro-organism(s) present in the sample from the differential hybridization signals obtained in step (iv).

30

The part of the ITS mentioned in the step of amplification, is a polynucleotide comprising the target sequence, or the target sequence itself, the target sequence consisting of SEQ ID NO 1 or 2, or of the RNA form of said SEQ ID NO 1 or 2

wherein T is replaced by U, or of the complementary form of said SEQ ID NO 1 or 2, or of any homologues, or of a fragment of at least 20 contiguous nucleotides thereof.

Preferentially, the present invention provides for a method as described above
5 wherein at least 2 micro-organisms are detected simultaneously.

A set of probes as described in step (iii) comprises at least two, three, four, five, six, seven, eight, nine or more probes of the invention, or equivalents thereof.

In a preferred method of the invention, set of probes as described in step (iii) comprises at least two probes.

10 Preferred probes are polynucleotides of SEQ ID NO 1 to 70 and their homologues.

The present invention also provides for a method as described above, wherein the probes as specified in step (iii) are combined with at least one other probe, preferentially also from the 16S-23S rRNA spacer region, enabling the simultaneous detection of
15 different pathogenic bacteria liable to be present in the same sample.

Preferred probes are designed for attaining optimal performance under the same
hybridization conditions so that they can be used in sets for simultaneous hybridization;
this highly increases the usability of these probes and results in a significant gain in time
20 and labor.

A kit containing any of the polynucleotides of the present invention is also an object of the invention.

25 A kit of the invention comprise the following components:

- at least one polynucleotide hybridizing to the target sequence consisting of SEQ ID NO 1 or 2, to the RNA form of said SEQ ID NO 1 or 2 wherein T is replaced by U, to the complementary form of said SEQ ID NO 1 or 2, or to any of their homologues;
- 30 - a hybridization buffer, or components necessary for producing said buffer.

A preferred kit comprises

- at least one set of two HybProbes hybridizing, adjacent to each other with less than 25 nucleotides, preferably less than 5 nucleotides, to the target sequence consisting

of SEQ ID NO 1 or 2, to the RNA form of said SEQ ID NO 1 or 2 wherein T is replaced by U, to the complementary form of said SEQ ID NO 1 or 2, or to any of their homologues;

- a hybridization buffer, or components necessary for producing said buffer.

Table1

SEQ IDs	Use	References	Length	Sequence
SEQ ID NO 1				TTTGACATTGAAXACTAGATAAGTAAGTAXAATATAGATTTACCAYGCAAAACCGAGTGAATA AAGYGTITTTAAZAAGCWTGAATTCATAVAAATAATCGZTAGZGTCGAYAGAACACTCACAA GUTTAATAACWSGT
SEQ ID NO 2				TTTGACATTGAAAACCTAGATAAGTAAGTAAAAATATAGATTTACCAAGCAAAACCGAGTGAATA AAGAGTTTAAATAAGCTTGAATTCATAAGAAATAATCGTAGTGTTCGAAAGAACACTCACAA GATTATAACGCGT
SEQ ID NO 3	CO	RStau6WTcompl.1b	63	ACGCTCACATACGGCTTCGTTTTCATTATTTAAATGCTCATTACATAAGTAACTCTGCTT
SEQ ID NO 4	CO	WToligoHP5	69	CTTATGAATTCAGGCTTATTTAAACTCTTTATTCACTCGGTTTTCGTTGGTAAATCTATATTT ACT
SEQ ID NO 5	CO	WToligoHP6	70	CGAACACTAGCGATTATTTCTATGAATTCAGGCTTATTTAAACTCTTTATTCACTCGGTTTTC CTTGG
SEQ ID NO 6	CO	RStau7WTcompl.1b	81	TTATAAGTCAACCGTTAACATGAAGTACGTTCTTTTATAAAAGATTAAACGCGTTATTAATCT TGTGAGTGTCTTTC
SEQ ID NO 7	CO	RStau6WTcompl.1a	83	TATAAGTCAACGCTCACATACGGCTTCGTTTTCATTATTTAAATGCTCATTACATAAGTAA CTCTGCTTTAAATAATT
SEQ ID NO 8	CO	RStauHP1WTcompl.1	88	CTTATTTAAACTCTTTATTCACTCGGTTTTCGTTGGTAAATCTATATTTACTTACTATCTAG TTTCAATGTACAAATAATGGT
SEQ ID NO 9	CO	WToligo-StauHP4	88	ATTTAAACTCTTTATTCACTCGGTTTTCGTTGGTAAATCTATATTTACTTACTTATCTAGTTT CAATGTACAATTTCTTTTAG
SEQ ID NO 10	CO	RStauHP2WTcompl.1	89	GTGAGTGTCTTTTCGAACACTAGCGATTATTTCTATGAATTCAGGCTTATTTAAACTCTTTATT CACTCGGTTTTCGTTGGTAAAT
SEQ ID NO 11	CO	RStauHP2cWtcompl.1	89	ATTTACCAAGCAAAACCGAGTGAATAAGAGTTTAAATAAGCTTGAATTCATAAGAAATAATC GCTAGTGTTCGAAAGAACACTCAC
SEQ ID NO 12	CO	RStauHP3WTcompl.1	91	TTTAAACGCGTTATTAATCTTGTGAGTGTCTTCGAACACTAGCGATTATTTCTATGAATTC AGCTTATTTAAACTCTTTATTCAT
SEQ ID NO 13	CO	RStau7WTcompl.1a	101	TCCACCATTTTATAAGTCAACGTTAACATGAAGTACGTTCTTTTATAAAAGATTAAACGCG GTTATTAATCTTGTGAGTGTCTTCGAACACTAGC
SEQ ID NO 14	HP	RStaphSP-5LC8.1	21	AGATTTTACCAAGCAAAACCG
SEQ ID NO 15	HP	RStauHP5-3FL1	21	AGATTTTACCAAGCAAAACCG
SEQ ID NO 16	HP	RStauHP6a-5LC6.1	21	AAGCTTGAATTCATAAGAAAT
SEQ ID NO 17	HP	RStauHP6-3FL1	23	CCGAGTGAATAAAGAGTTTAA
SEQ ID NO 18	HP	RStauHP4-5LC6.1	24	CCAAGCAAAACCGAGTGAATAAAG
SEQ ID NO 19	HP	RStauHP6b-5LC6.1	24	GCTTGAATTCATAAGAAATAATCG
SEQ ID NO 20	HP	RStauHP5b-5LC6.1	25	GAATAAAGAGTTTAAATAAGCTTG
SEQ ID NO 21	HP	RStauHP5a-5LC6.1	27	GTGAATAAAGAGTTTAAATAAGCTTG
SEQ ID NO 22	HP	RStauHP9-5LC6	27	AAGCTTGAATTCATAAGAAATAATCGC
SEQ ID NO 23	HP	RStaphHP8-5LCR6	28	CTTGGTAAATCTTACTTACTATCTAG
SEQ ID NO 24	HP	RStau63FL1	29	AAGCAGAGTTTACTTATGTAAATGAGCAT
SEQ ID NO 25	HP	RStauHP1-5LCR64.1	29	TACCAAGCAAAACCGAGTGAATAAAGAGT
SEQ ID NO 26	HP	RStaphSP-3FL1	29	TTTGACATTGAAAACCTAGATAAGTAAGT
SEQ ID NO 27	HP	RStauHP7-3FL1	29	GCAAAACCGAGTGAATAAAGAGTTTAA
SEQ ID NO 28	HP	RStauHP7a-5LCR6	30	AAGCTTGAATTCATAAGAAATAATCGCTAG
SEQ ID NO 29	HP	RStauHP9-3FL1	30	AGCAAAACCGAGTGAATAAAGAGTTTAA
SEQ ID NO 30	HP	RStauHP4-3FL1	31	ATTGTACATTGAAAACCTAGATAAGTAAGTAA
SEQ ID NO 31	HP	RStauHP3-4LC6.1	32	AGTGTTCGAAAGAACACTCACAGATTAAATA
SEQ ID NO 32	HP	RStauHP8-3FL1	32	AGCTTATTTAAACTCTTTATTCACTCGGTTT
SEQ ID NO 33	HP	RStau65LC6.1	33	TAAAAATATGAAAACGAAGCCGATGTGAGCGT
SEQ ID NO 34	HP	RStau65LC7.1	33	TAAAAATATGAAAACGAAGCCGATGTGAGCGT
SEQ ID NO 35	HP	RStauHP2-4LC6.1	33	GAATTCATAAGAAATAATCGCTAGTGTTCGAA
SEQ ID NO 36	HP	RStauHP2c-3FL1	33	TTTCGAACACTAGCGATTATTTCTTATGAATTC
SEQ ID NO 37	HP	RStauHP2-3FL1	34	GCAAAACCGAGTGAATAAAGAGTTTAAATAAGC
SEQ ID NO 38	HP	RStauHP2c-4LC6.1	34	GCTTATTTAAACTCTTTATTCACTCGGTTTTCG
SEQ ID NO 39	HP	RStauHP8-5LCR6	36	GCTTGGTAAATCTATATTTACTTACTTATCTAGT
SEQ ID NO 40	HP	RStauHP1-3FL1	38	GTACATTGAAAACCTAGATAAGTAAGTAAATATAGATT
SEQ ID NO 41	HP	RStauHP3-3FL1	38	GAGTTTTAAATAAGCTTGAATTCATAAGAAATAATCGC
SEQ ID NO 42	HP	RStau73FL1	39	GAAAGAACACTCACAGATTAAATAACGCGTTTAAATCTT
SEQ ID NO 43	HP	RStau75LC6.1	41	TTATAAAAGAACGTAACCTCATGTAAACGTTTGACTTATAA
SEQ ID NO 44	HP	RStau75LC7.1	41	TTATAAAAGAACGTAACCTCATGTAAACGTTTGACTTATAA
SEQ ID NO 45	Pr	RStauFP18.1	17	CTTCAGAAGATGCGGAA
SEQ ID NO 46	Pr	RStauFP22.1	18	TTTCGAACACTAGCGATT
SEQ ID NO 47	Pr		20	GTTATTAUCTTGTGAGTGTT
SEQ ID NO 48	Pr		20	ACSGTTTATTAUCTTGTGAG
SEQ ID NO 49	Pr	RStauFP19.1	19	CTTCAGAAGATGCGGAA
SEQ ID NO 50	Pr	RStauFP20.1	19	TTCTTCAGAAGATGCGGAA
SEQ ID NO 51	Pr	Staph-P26rev	19	TTCTTCGAACACTAGCGA
SEQ ID NO 52	Pr	StauP01	20	ACCAAGCAAAACCGAGTGAA
SEQ ID NO 53	Pr	Staph-P25rev	20	GTTCTTCGAACACTAGCGA
SEQ ID NO 54	Pr		20	SGGTATTAUCTTGTGAGTG
SEQ ID NO 55	Pr	StaphP28rev	20	CGCGTTATTAATCTGTGAG

SEQ IDs	Use	References	Length	Sequence
SEQ ID NO 56	Pr		20	CATTGAAXACTAGATAAGTA
SEQ ID NO 57	Pr		20	GTTATTAAUCTTGTGAGTGTT
SEQ ID NO 58	Pr	SlaphP31rev	22	ACGCGTTATTAACTTGTGAGT
SEQ ID NO 59	Pr	RstauRP21.1	23	CGAACACTAGCGATTATTCTTA
SEQ ID NO 60	Pr	RStauRP23.1	23	ACTAGCGATTATTCTTATGAAT
SEQ ID NO 61	Pr	SlaphP34	23	CATATTGTATTACGTTTGAATG
SEQ ID NO 62	Pr		18	SXGTTATTAAUCTTGTGAG
SEQ ID NO 63	Pr	SlaphP32	24	CATATTGTATTCAGTTTGAATGC
SEQ ID NO 64	Pr	SlaphP33	24	CATATTGTATTCAGTTTGAATGT
SEQ ID NO 65	Pr	StauP11	25	CATAAGAAATATCGCTAGTGTTG
SEQ ID NO 66	Pr	Slaph-P24	25	GACATATTGTATTCAGTTTGAATG
SEQ ID NO 67	Pr	SlaphP29	26	GAAAACTAGATAAGTAAGTAAATAT
SEQ ID NO 68	Pr	SlaphP30	26	TGTACATTGAAAAGTATAGTAAG
SEQ ID NO 69	Pr		24	TTTGTACATTGAAXACTAGATAAG
SEQ ID NO 70	Pr		24	ACSXGTTATTAAUCTTGTGAGTGTT

Use is in fact preferred use as HP =HybProbe, Pr = primer, CO = complementary polynucleotides.

5 S = A, T, C or G

U = A or T

V = G or C

W = T or G

X = A or C

10 Y = A or G

Z = T or C

- <SEQ ID NO:1;DNA;staphylococcus>
TTTGACATTGAAMACTAGATAAGTAAGTAMAATATAGATTTTACCARGCAAAACCGAGT
GAATAAAAGRGTTTTTRAAYAAGCKTGAATTCATAASAAATAATCGYTAGYGTTTCGARAGAA
CACTACAAGWTTAATAACKNGT
- 5 <SEQ ID NO:2;DNA;staphylococcus>
TTTGACATTGAAAACCTAGATAAGTAAAGTAAATATAGATTTTACCAAGCAAAACCGAGT
GAATAAAGAGTTTTAAATAAGCTTGAATTCATAAGAAATAATCGCTAGTGTTTCGAAAGAA
CACTACAAGATTAATAACGCGT
- 10 <SEQ ID NO:3;DNA;staphylococcus>
ACGCTCACATACGGCTTCGTTTTTCATTATTTAAATGCTCATTTACATAAGTAAACTCTGCT
T
- 15 <SEQ ID NO:4;DNA;staphylococcus>
CTTATGAATTCAAGCTTATTTAAACTCTTTATTCACCTCGGTTTTGCTTGGTAAAATCTATA
TTTTACT
- <SEQ ID NO:5;DNA;staphylococcus>
20 CGAACACTAGCGATTATTTCTTATGAATTCAAGCTTATTTAAACTCTTTATTCACCTCGGTT
TTGCTTGG
- <SEQ ID NO:6;DNA;staphylococcus>
25 TTATAAGTCAAACGTTAACATGAAGTTACGTTCTTTTATAAAAAGATTTAAACGCGTTATT
AATCTTGTGAGTGTTCTTTC
- <SEQ ID NO:7;DNA;staphylococcus>
TATAAGTCAAACGCTCACATACGGCTTCGTTTTTCATTATTTAAATGCTCATTTACATAAGT
AAACTCTGCTTTAAATAATT
- 30 <SEQ ID NO:8;DNA;staphylococcus>
CTTATTTAAACTCTTTATTCACCTCGGTTTTGCTTGGTAAAATCTATATTTACTTACTTATC
TAGTTTTCAATGTACAAATAATGGT
- 35 <SEQ ID NO:9;DNA;staphylococcus>
ATTTAAACTCTTTATTCACCTCGGTTTTGCTTGGTAAAATCTATATTTACTTACTTATCTA
GTTTTCAATGTACAATTCTTTTTAG
- <SEQ ID NO:10;DNA;staphylococcus>
40 GTGAGTGTTCTTTTCGAACACTAGCGATTATTTCTTATGAATTCAAGCTTATTTAAACTCTT
TATTCACCTCGGTTTTGCTTGGTAAAAT
- <SEQ ID NO:11;DNA;staphylococcus>
45 ATTTTACCAAGCAAAACCGAGTGAATAAAGAGTTTTAAATAAGCTTGAATTCATAAGAAA
TAATCGCTAGTGTTTCGAAAGAACAACCTCAC
- <SEQ ID NO:12;DNA;staphylococcus>
TTTAAACGCGTTATTAATCTTGTGAGTGTTCTTTTCGAACACTAGCGATTATTTCTTATGAAT
TCAAGCTTATTTAAACTCTTTATTCAC
- 50 <SEQ ID NO:13;DNA;staphylococcus>
TCCACCATTTTTATAAGTCAAACGTTAACATGAAGTTACGTTCTTTTATAAAAAGATTTAA
ACGCGTTATTAATCTTGTGAGTGTTCTTTTCGAACACTAGC
- 55 <SEQ ID NO:14;DNA;staphylococcus>
AGATTTTACCAAGCAAAACCG
- <SEQ ID NO:15;DNA;staphylococcus>
60 AGATTTTACCAAGCAAAACCG

- <SEQ ID NO:16;DNA;staphylococcus>
AAGCTTGAATTCATAAGAAAT
- 5 <SEQ ID NO:17;DNA;staphylococcus>
CCGAGTGAATAAAGAGTTTAAA
- <SEQ ID NO:18;DNA;staphylococcus>
CCAAGCAAAACCGAGTGAATAAAG
- 10 <SEQ ID NO:19;DNA;staphylococcus>
GCTTGAATTCATAAGAAATAATCG
- <SEQ ID NO:20;DNA;staphylococcus>
15 GAATAAAGAGTTTTAAATAAGCTTG
- <SEQ ID NO:21;DNA;staphylococcus>
GTGAATAAAGAGTTTTAAATAAGCTTG
- 20 <SEQ ID NO:22;DNA;staphylococcus>
AAGCTTGAATTCATAAGAAATAATCGC
- <SEQ ID NO:23;DNA;staphylococcus>
CTTGGTAAATCTTACTTACTTATCTAG
- 25 <SEQ ID NO:24;DNA;staphylococcus>
AAGCAGAGTTTACTTATGTAAATGAGCAT
- <SEQ ID NO:25;DNA;staphylococcus>
30 TACCAAGCAAAACCGAGTGAATAAAGAGT
- <SEQ ID NO:26;DNA;staphylococcus>
TTTGTACATTGAAAAGTAGATAAGTAAGT
- 35 <SEQ ID NO:27;DNA;staphylococcus>
GCAAAACCGAGTGAATAAAGAGTTTAAA
- <SEQ ID NO:28;DNA;staphylococcus>
AAGCTTGAATTCATAAGAAATAATCGCTAG
- 40 <SEQ ID NO:29;DNA;staphylococcus>
AGCAAAACCGAGTGAATAAACAGTTTTAAA
- <SEQ ID NO:30;DNA;staphylococcus>
45 ATTGTACATTGAAAAGTAGATAAGTAAGTAA
- <SEQ ID NO:31;DNA;staphylococcus>
AGTGTTTCGAAAGAACAACCTCACAAGATTAATAA
- 50 <SEQ ID NO:32;DNA;staphylococcus>
AGCTTATTTAAACTCTTTATTCACCTCGGTTT
- <SEQ ID NO:33;DNA;staphylococcus>
55 TAAAATAATGAAAACGAAGCCGTATGTGAGCGT
- <SEQ ID NO:34;DNA;staphylococcus>
TAAAATAATGAAAACGAAGCCGTATGTGAGCGT
- 60 <SEQ ID NO:35;DNA;staphylococcus>

- GAATTCATAAGAAATAATCGCTAGTGTTTCGAAA
- <SEQ ID NO:36;DNA;staphylococcus>
TTTCGAACACTAGCGATTATTTCTTATGAATTC
- 5 <SEQ ID NO:37;DNA;staphylococcus>
GCAAAACCGAGTGAATAAAGAGTTTTAAATAAGC
- <SEQ ID NO:38;DNA;staphylococcus>
10 GCTTATTTAAACTCTTTATTCACCTCGGTTTTGC
- <SEQ ID NO:39;DNA;staphylococcus>
GCTTGGTAAAACTATATTTACTTACTTATCTAGT
- 15 <SEQ ID NO:40;DNA;staphylococcus>
GTACATTGAAAACCTAGATAAGTAAGTAAATATAGATT
- <SEQ ID NO:41;DNA;staphylococcus>
GAGTTTTAAATAAGCTTGAATTCATAAGAAATAATCGC
- 20 <SEQ ID NO:42;DNA;staphylococcus>
GAAAGAACACTACAAGATTAATAACGCGTTTAAATCTT
- <SEQ ID NO:43;DNA;staphylococcus>
25 TTATAAAAGAACGTAACTTCATGTTAACGTTTGACTTATAA
- <SEQ ID NO:44;DNA;staphylococcus>
TTATAAAAGAACGTAACTTCATGTTAACGTTTGACTTATAA
- 30 <SEQ ID NO:45;DNA;staphylococcus>
CTTCAGAAGATGCGGAA
- <SEQ ID NO:46;DNA;staphylococcus>
TTTCGAACACTAGCGATT
- 35 <SEQ ID NO:47;DNA;staphylococcus>
GTTATTAAWCTTGTGAGTGTT
- <SEQ ID NO:48;DNA;staphylococcus>
40 ACNMGTTATTAAWCTTGTGAG
- <SEQ ID NO:49;DNA;staphylococcus>
CTTCAGAAGATGCGGAATA
- 45 <SEQ ID NO:50;DNA;staphylococcus>
TTCTTCAGAAGATGCGGAA
- <SEQ ID NO:51;DNA;staphylococcus>
TTCTTTCGAACACTAGCGA
- 50 <SEQ ID NO:52;DNA;staphylococcus>
ACCAAGCAAAACCGAGTGAA
- <SEQ ID NO:53;DNA;staphylococcus>
55 GTTCTTTCGAACACTAGCGA
- <SEQ ID NO:54;DNA;staphylococcus>
NMGTTATTAAWCTTGTGAGTG
- 60 <SEQ ID NO:55;DNA;staphylococcus>

CGCGTTATTAATCTTGTGAG

<SEQ ID NO:56;DNA;staphylococcus>
CATTGAAMACTAGATAAGTA

5

<SEQ ID NO:57;DNA;staphylococcus>
GTTATTAAWCTTGTGAGTGTT

10

<SEQ ID NO:58;DNA;staphylococcus>
ACGCGTTATTAATCTTGTGAGT

<SEQ ID NO:59;DNA;staphylococcus>
CGAACACTAGCGATTATTCTTA

15

<SEQ ID NO:60;DNA;staphylococcus>
ACTAGCGATTATTCTTATGAAT

<SEQ ID NO:61;DNA;staphylococcus>
CATATTGTATTCAGTTTGAATG

20

<SEQ ID NO:62;DNA;staphylococcus>
NMGTTATTAAWCTTGTGAG

<SEQ ID NO:63;DNA;staphylococcus>
CATATTGTATTCAGTTTGAATGC

25

<SEQ ID NO:64;DNA;staphylococcus>
CATATTGTATTCAGTTTGAATGT

30

<SEQ ID NO:65;DNA;staphylococcus>
CATAAGAAATAATCGCTAGTGTTCG

<SEQ ID NO:66;DNA;staphylococcus>
GACATATTGTATTCAGTTTGAATG

35

<SEQ ID NO:67;DNA;staphylococcus>
GAAAAC TAGATAAGTAAGTAAAATAT

<SEQ ID NO:68;DNA;staphylococcus>
TGTACATTGAAAAC TAGATAAGTAAG

40

<SEQ ID NO:69;DNA;staphylococcus>
TTTGTACATTGAAMACTAGATAAG

45

<SEQ ID NO:70;DNA;staphylococcus>
ACNMGTTATTAAWCTTGTGAGTGTT

Table 2

SEQ ID s NO	62 (RP)	57 (RP)	59 (RP)	46 (RP)	60 (RP)	53 (RP)	51 (RP)	54 (RP)	55 (RP)	58 (RP)
52 (FP)	1									
65 (FP)		2								
45 (FP)			3	4	5					
49 (FP)			6	7	8					
50 (FP)			9	10	11					
66 (FP)	32					12	13	14	15	16
67 (FP)									17	18
68 (FP)	33								19	20
63 (FP)	21								22	23
64 (FP)	24									
61 (FP)	25		31			26		27	28	29

FP/RP = Forward primers / Reverse primers

= Preferred primer combinations

Table 3

<u>SEQ IDs</u> <u>NO</u>	<u>Hybprobe</u>	<u>Primercombination</u>	<u>Protocol PCR</u>	<u>Performance in the</u> <u>particular conditions of</u> <u>the examples</u>
24 / 33	RStau63Fl.1 / RStau65LC6.1(LCR640)	1, 2	$t_a = 60^\circ\text{C}$	+
24 / 34	RStau63Fl.1 / RStau65LC7.1 (LCR705)	1	$t_a = 60^\circ\text{C}$	+
42 / 43	RStau73Fl.1 / RStau75LC6.1 (LCR640)	1, 2	$t_a = 60^\circ\text{C}$	+
42 / 44	RStau73Fl.1 / RStau75LC7.1 (LCR705)	1	$t_a = 60^\circ\text{C}$	+
40 / 25	RStauHP1-3Fl.1 / RStauHP1- 5LCR64.1	3, 4, 5, 6, 7, 8, 9, 10, 11, 27, 29, 31	$t_a = 55^\circ\text{C}$	++
37 / 35	RStauHP2-3Fl.1 / RStauHP2-LC6.1	12, 13, 15, 16, 17, 18, 19, 20, 22, 23, 28	$t_a = 55^\circ\text{C}$	++
36 / 38	RStauHP2c-3Fl.1 / RStauHP2c-LC6.1	14, 20, 33	$t_a = 55^\circ\text{C}$	+
41 / 31	RStauHP3-3Fl.1 / RStauHP3-LC6.1	14, 25, 33	$t_a = 55^\circ\text{C}$	++
26 / 14	RStaphSP-3Fl.1 / RStaphSP-5LC6.1	6, 12, 14, 21, 24, 25, 27, 29, 32	$t_a = 55^\circ\text{C}$	+
30 / 18	RStauHP4-3Fl.1 / RStauHP4-5LC6.1	29	$t_a = 55^\circ\text{C}$	+
15 / 21	RStauHP5-3Fl.1 / RStauHP5a-5LC6.1	20	$t_a = 55^\circ\text{C}$	+
15 / 20	RStauHP5-3Fl.1 / RStauHP5b-5LC6.1	20	$t_a = 55^\circ\text{C}$	+
17 / 16	RStauHP6-3Fl.1 / RStauHP6a-5LC6.1	20	$t_a = 55^\circ\text{C}$	+
17 / 19	RStauHP6-3Fl.1 / RStauHP6b-5LC6.1	20	$t_a = 50^\circ\text{C}$	+++
27 / 28	RStauHP7-3Fl.1 / RStauHP7a-5LCR6	20	$t_a = 50^\circ\text{C}$	+
32 / 39 / 23	RStauHP8.3Fl.1 / RStauHP8.5LCR6 / RStepHP8.5LCR6	29	$t_a = 55^\circ\text{C}$	+
29 / 22	RStauHP9-3Fl.1 / RStauHP9-5LC6	20	$t_a = 50^\circ\text{C}$	+

* The primer pairs in bold performed the best.

Table4

<i>Staphylococci</i>				
Species (32)	Subspecies (15)	Clinical relevance	Coagulase reaction	Examined in this study
<i>S. aureus</i>	<i>aureus</i>	++++	+	+
	<i>anearobius</i>	++++	+	+
<i>S. epidermidis</i>		+++	-	+
<i>S. capitis</i>	<i>capitis</i>	+	-	+
	<i>ureolyticus</i>	+	-	+
<i>S. caprae</i>		+	-	+
<i>S. saccharolyticus</i>		+	-	+
<i>S. warneri</i>		+	-	+
<i>S. pasteurii</i>		+	-	+
<i>S. haemolyticus</i>		+++	-	+
<i>S. hominis</i>	<i>hominis</i>	+	-	+
	<i>novobiosepticum</i>	+	-	-
<i>S. lugdunensis</i>		+	-	+
<i>S. schleiferi</i>	<i>schleiferi</i>	+	-	+
	<i>coagulans</i>	+	+	+
<i>S. muscae</i>		-	-	-
<i>S. auricularis</i>		+	-	+
<i>S. saprophyticus</i>	<i>saprophyticus</i>	++	-	+
	<i>bovis</i>	+	-	-
<i>S. cohnii</i>	<i>cohnii</i>	+	-	+
	<i>urealyticum</i>	+	-	+
<i>S. xylosus</i>		- (+)	-	+
<i>S. kloosii</i>		-	-	-
<i>S. equorum</i>		-	-	-
<i>S. arletae</i>		-	-	-
<i>S. gallinarum</i>		-	-	-
<i>S. simulans</i>		+	-	+
<i>S. carnosus</i>		-	-	-
<i>S. piscifermentans</i>		-	-	-
<i>S. felix</i>		-	-	-
<i>S. lutrae</i>		-	+	-
<i>S. intermedius</i>		-	+	-
<i>S. delphini</i>		-	+	-
<i>S. hyicus</i>		-	(+)	+
<i>S. chromogenes</i>		-	-	+
<i>S. sciuri</i>	<i>sciuri</i>	-	-	+
	<i>carnaticus</i>	-	-	-
	<i>rodentium</i>	-	-	-
<i>S. lentus</i>		-	-	-
<i>S. vitulus</i>		-	-	-

EXAMPLES

The method used in the examples is a method for the detection of *Staphylococci*,
5 in particular *S. aureus*, *S. epidermidis* and *S. haemolyticus*, using the HybProbes of SEQ
ID NO 17 and 19, in combination with a *Staphylococcus*-genus primer set of SEQ ID 58
and 68.

The HybProbes are labeled: SEQ ID NO 17 in 3' with fluorescein and SEQ ID NO
19 in 5' with LCred640.

10 In total, 63 *S. aureus* isolates (including one representative of the *S. aureus* subs.
anaerobius), 48 *S. epidermidis* isolates and 16 *S. haemolyticus* isolates from different
geographical origins were investigated.

If the isolates gave not the expected results, the gDNA was retyped by t-RNA PCR
(Vaneechoutte, M. et al., 1998, Int. J. Syst. Bacteriol. (48) 127-139) and/or the culture
15 was retyped by ApiSTAPH (Biomérieux).

The instrumentation is the LightCycler™ (version 1.2) provided with the adequate
software (LC-software version 3.5) enabling a Real-Time fluorescence PCR detection.

20

Example 1: Preparation of the samples to be tested**1/. DNA from pure cultures**

For extracting the DNA from pure cultures, different purification methods can be
25 used:

- Lysis with lysostaphin (5µg/µl) for 1 h at 37°C and purification with the
QIAamp blood DNA isolation kit (Qiagen)
- The method of Pitcher et al. (1989)
- The MagNAPure LC DNA isolation Kit III (Bacteria, Fungi) on the
30 MagNAPure instrument. Bacterial cells grown O/N on LB plates or slants
were suspended in 100 to 1000 µl TE pH8 for storage at -20°C. 2 µl to 20 µl
was used for extraction according to the manufacturer's recommendations.

- QIAamp DNA mini kit (catalog no. 51306 – QIAGEN). The culture was pre-treated enzymatically using lysozyme and lysostaphin.

5 2/. DNA from positive blood culture bottles

Blood samples were inoculated in aerobe blood culture bottles (BacT/ALERT FA) and incubated in a BacT/Alert 3D system (Organon Teknika) at 37°C until positive. Positivity was monitored by a color change from dark green to yellow.

Aliquots (1.5ml) of the blood cultures were frozen at -70°C until use.

- 10 Genomic DNA was prepared as described in the pack insert of the MPLC DNA Isolation Kit III. As recommended for Organon Teknika blood culture bottles, prior to PCR the eluate was centrifuged 10 sec at 14000rpm to spin down the extracted carbon particles.

15

Example 2: LightCycler (LC) protocol

Following the instructions of the manufacturer of the kit LC-FastStart DNA Master Hybridization Probes (cat. No 3 003 248 or No 2 239 272):

- 20 - any sample material suitable for PCR in terms of purity, concentration, and absence of inhibitors can be used ;
- the primers should be at a final concentration of 0,3 to 1 μ M each ;
 - the HybProbes at a final concentration of 0,2 μ M each, or double ;
 - the concentration of MgCl₂ should be optimized, and may vary from 1 to
- 25 5mM;
- and a negative control should be run.

- 30 The amplification and melting conditions are described herein after. The LC software version 3.5 was used. The quantification settings were F2/back F1 (samples). For the baseline adjustment the arithmetic mode was used. The crossing point (Ct) calculation was based on the second derivative maximum. The calculation method for the melting peak was polynomial. The peak area was used to calculate the Tm.

Amplification and melting curve program:

	Temp.(°C)	Hold time	Slope (°C/sec.)	Acquisition mode
Denaturation	95	10 min	20	None
45x { Cycles	95	10 sec	20	None
	50	15 sec	20	SINGLE
	72	10 sec	20	None
Melting	95	60 sec	20	None
	40	60 sec	20	None
	80	0 sec	0.1	CONTINUOUS
Cooling	40	0 sec	20	None

5

Example 3: Results on purified DNA, inclusivity and cross reactivity tests

1/ Inclusivity

All *S. aureus* isolates examined (n=63) were successfully amplified (Ct range
 10 17.25 - 33.51) and gave one uniform melting peak with a mean T_m of 53.13°C ± 0.52°C
 whatever the geographical or specimen origin.

It has to be noted that the subspecies *S. aureus* subs. *anaerobius* fell together with
 the *S. aureus* species.

15 All isolates received as *S. epidermidis* (n=48) and *S. haemolyticus* (n=16) could be
 detected using the melting curve. Quantification curves usually were not observed. The
 average T_m value for *S. epidermidis* isolates was 44.55°C ± 0.21°C and for *S.*
haemolyticus isolates it was 44.96°C ± 0.24°C. There were no differences observed
 among isolates from different geographical or specimen origin.

20

All isolates received as *S. aureus* or *S. haemolyticus* reacted as expected. However,
 deviating results were obtained for five *S. epidermidis* isolates.

Two of these, one from UK and one from Italy, were afterwards identified as *S.*
hominis. Another *S. epidermidis* isolate from UK showed a T_m value of 49.02°C and
 25 could afterwards only be determined as a *Staphylococcus* species (not *S. aureus*, *S.*
epidermidis or *S. haemolyticus*). The UK isolate was afterwards retyped as a *S.*

haemolyticus. And one isolate from Spain producing an uncharacteristic small melting peak was retyped as *S. intermedius* or *S. chromogenes*.

2/. Cross-reactivity

5 More than 50 different bacterial species were tested (*Mycobacteria*, *Pseudomonas*, *Streptococci*, etc) and also few fungi. None of the tested organisms generated quantification curves or melting peaks with the assay.

3/. Conclusion:

10 All *S. aureus* isolates investigated are detected (100% sensitivity) and could be unequivocally distinguished from all other isolates studied.

Also for *S. epidermidis* and *S. haemolyticus* the sensitivity was 100%.

Therefore using this particular set of HybProbes of SEQ IDs NO 17 and 19, both species are detected without being differentiated from each other or from other CoNS.

15 Unwanted cross-reactivities, with organisms from the pathogens tested were not observed.

Summary of the sensitivity and specificity tests.

Taxon	# of strains tested	# of strains with		
		Peak at 53°C	Peak below 49°C	No peak
<i>S.aureus</i>	63	63	0	0
<i>S.epidermidis</i>	43	0	43	0
<i>S.haemolyticus</i>	16	0	16	0
Cross-check list	55	0	0	55
Human DNA	1	0	0	1

20

Example 4: Results on blood cultures

In total aliquots of 16 positive blood culture bottles inoculated with patient-
25 samples were tested.

From the five bottles that grew *S. aureus*, 4 could be identified as such. One showed no peak. Indeed, the DNA of the sample was retyped as *S. schleiferi*.

Out of the ten bottles found positive for *S. epidermidis*, 7 showed a peak at 44-45°C – as expected – while three showed peaks shifted to 47°C. After retyping the DNA of the deviating samples, they were identified as *S. hominis*.

One bottle, positive for *S. haemolyticus*, produced the expected melting peak at
5 44.59°C

Example 5: Detection of other

10 In total 51 CoNS-strains, other than *S. epidermidis* and *S. haemolyticus*, have been studied. The results are summarized in the table below.

Amplification was observed on gel for all strains of all species except two (*S. schleiferi* and *S. sciuri*) for which the results are ambiguous (see also below). Some CoNS produced a growth curve (amplification), with corresponding Ct value, but the
15 endpoint fluorescence was very low (0.004 or less) as compared to the values obtained for *S. aureus* (e.g. 0.03 for 10³ copies).

Most species examined were detectable by melting curve analysis, with the exception of *S. chromogenes*, *S. hyicus* and *S. simulans*.

The results for *S. schleiferi* and *S. sciuri* are obscure; for both species melting
20 peaks were observed but could not be discriminated with certainty from a potential contaminating agent. Results obtained with universal primers suggest that both species will presumably not be detectable; there was clear prove of amplification on gel, but the HybProbes did not generate a signal in the LC.

The T_m range observed for CoNS is 39 to 48°C. This is clearly different from that
25 of *S. aureus* but perfectly overlapping the range of *S. epidermidis* and *S. haemolyticus*.

Summary of the LC results obtained for strains of the *Staphylococcus* species examined.

Species	# strains tested	Amplicon on gel (+/-140 bp)	Ct detection	Range Tm(°C) melting peak	Melting peak detection
<i>S. aureus</i>	70	+	Yes	52-54	Yes
<i>S. epidermidis</i>	12	+	No	43-45	Yes
<i>S. haemolyticus</i>	21	+	No	44-46 (47)	Yes
<i>S. auricularis</i>	3	+	No	39-40	(Yes)
<i>S. capitis</i>	4	+	No	46-47	Yes
<i>S. caprae</i>	2	+	No	45-46	Yes
<i>S. chromogenes</i>	1	+	No	-	No
<i>S. cohnii</i>	3	+	(Yes)	46-48	Yes
<i>S. hominis</i>	6	+	No	46-48	Yes
<i>S. hyicus</i>	1	+	No	-	No
<i>S. lugdunensis</i>	12	+	(Yes)	46-48	Yes
<i>S. pasteurii</i>	1	+	(Yes)	47	Yes
<i>S. saccharolyticus</i>	1	+	No	45	Yes
<i>S. saprophyticus</i>	5	+	No	42-43	Yes
<i>S. schleiferi</i>	4	?	No	?	?
<i>S. sciuri</i>	1	?	No	?	?
<i>S. simulans</i>	1	+	No	-	No
<i>S. warneri</i>	5	+	(Yes)	(42) 46-47	Yes
<i>S. xylosus</i>	1	+	No	42	Yes
Total	154				

CLAIMS

1. An isolated nucleic acid molecule consisting of SEQ ID NO 1, its complementary form, or RNA form thereof.

5

2. An isolated nucleic acid molecule consisting of SEQ ID NO 2, its complementary form, or RNA form thereof.

3. An isolated nucleic acid molecule that specifically hybridizes to SEQ ID NO 1 or 2, or to the RNA form of said SEQ ID NO 1 or 2 wherein T is replaced by U, or to the complementary form of said SEQ ID NO 1 or 2, or to a fragment of at least 20 contiguous nucleotides thereof, or to any of their homologues, for the detection and/or identification of *Staphylococcus* species, in particular of *S. aureus*.

4. An isolated nucleic acid molecule according to claim 3 consisting of a nucleic acid selected from the group consisting of SEQ IDs NO 14, 16 to 23, 25 to 32, 35 to 42, 51, 52, 53, 55, 58, 65, 67, 68, 69 and 70.

5. A set of two polynucleotide probes, said two probes hybridizing specifically to SEQ ID NO 1 or SEQ ID NO 2 or homologues, or to their RNA form wherein T is replaced by U, or to their complementary form, wherein there are no more than 25 nucleotides between said two probes.

6. A set of two polynucleotide probes according to claim 5 consisting of SEQ IDs NO 15 and 20, or SEQ IDs NO 15 and 21, or SEQ IDs NO 17 and 16, or SEQ IDs NO 17 and 19, or SEQ IDs NO 26 and 14, or SEQ IDs NO 27 and 28, or SEQ IDs NO 29 and 22, or SEQ IDs NO 32 and 39, or SEQ IDs NO 32 and 23, or SEQ IDs NO 30 and 18, or SEQ IDs NO 36 and 38, or SEQ IDs NO 37 and 35, or SEQ IDs NO 40 and 25, or SEQ IDs NO 41 and 31, or SEQ IDs NO 42 and 43.

30

7. A composition comprising at least one nucleic acid molecule according to any of claims 1 to 4 and/or a set of two polynucleotide probes according to claim 5 or claim 6.

8. Use of a nucleic acid molecule consisting of SEQ ID NO 1 or 2, or of the RNA form of said SEQ ID NO 1 or 2 wherein T is replaced by U, or of the complementary form of said SEQ ID NO 1 or 2, or of a fragment of at least 20 contiguous nucleotides thereof, or of any of their homologues, for the detection and/or
5 identification of *Staphylococcus* species, in particular of *S. aureus*.

9. A method for detecting or identifying *Staphylococcus* species using at least one nucleic acid molecule according to any of claims 1 to 4.

10. A method according to claim 9 for detection and/or identification of
10 *Staphylococcus* species in a sample comprising the steps of:

(i) if need be releasing, isolating and/or concentrating the polynucleic acids in the sample;

(ii) if need be amplifying the 16S-23S rRNA spacer region, or a fragment
15 comprising the target sequence, or the target sequence or a fragment thereof, with at least one suitable primer pair;

(iii) hybridizing the polynucleic acids of step (i) or (ii) with at least one polynucleotide probe that hybridizes to the target sequence,
wherein the target sequence of step (ii) and (iii) consists of SEQ ID NO 1 or 2 or
20 homologues thereof, or to their RNA form wherein T is replaced by U, or to their complementary form, or to a fragment of at least 20 contiguous nucleotides thereof,

(iv) detecting the hybrids formed, and

(v) interpreting the signal(s) obtained and inferring the presence of
Staphylococcus species and/or identifying the *Staphylococcus* species in the sample.

25

11. A method according to claim 10 wherein a suitable primer pair consists of any combination of a forward primer polynucleotide selected from the group consisting of SEQ ID NO 45, 49, 50, 52, 56, 61, 63, 64, 65, 66, 67, 68 and their homologues, and a reverse primer polynucleotide selected from the group consisting of SEQ ID NO 46, 47,
30 48, 51, 53, 54, 55, 57, 58, 59, 60, 62, and their homologues.

12. A method according to claim 10 or claim 11 wherein two polynucleotide probes are used.

13. A method according to claim 12 wherein the two polynucleotide probes hybridize to the target sequence adjacent to each other with less than 25 nucleotides in between.

5 14. A method according to claim 13 wherein the two polynucleotide probes consist of polynucleotides of SEQ IDs NO 15 and 20, or 15 and 21, or 17 and 16, or 17 and 19, or 26 and 14, or 27 and 28, or 29 and 22, or 32 and 39, or 32 and 23, or 30 and 18, or 36 and 38, or 37 and 35, or 40 and 25, or 41 and 31, or 42 and 43.

10 15. A kit for detection and/or identification of *Staphylococcus* species comprising the following components:

- at least one nucleic acid molecule according to any of claims 1 to 4 and/or a set of two polynucleotide probes according to claim 5 or 6.
- a hybridization buffer, or components necessary for producing said buffer.

15

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
24 June 2004 (24.06.2004)

PCT

(10) International Publication Number
WO 2004/052606 A3

(51) International Patent Classification⁷: **C12Q 1/68**

(21) International Application Number:
PCT/EP2003/013907

(22) International Filing Date: 8 December 2003 (08.12.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
02447247.4 6 December 2002 (06.12.2002) EP

(71) Applicants (for all designated States except US): **INNO-GENETICS N.V.** [BE/BE]; Technologiepark 6, B-9052 Ghent (BE). **ROCHE DIAGNOSTICS GMBH** [DE/DE]; Sandhofer Strasse 116, 68305 Mannheim (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CLAEYS, Sofie** [BE/BE]; Brieversweg 365, B-8310 Brugge-Sint-Kruis (BE). **JANNES, Geert** [BE/BE]; E. Vanhoorenbekelaan 23/1, B-3010 Leuven (BE). **HABERHAUSEN, Gerd** [DE/DE]; Kapellenwiese 35, 82377 Penzberg (DE). **EMRICH, Thomas** [DE/DE]; Blombergstrasse 7, 82393 Iffeldorf (DE). **VERDOODT, Lia** [BE/BE]; Wielewaalpark 10, B-9840 De Pinte (BE).

(74) Common Representative: **INNOGENETICS N.V.**; Intellectual Property Department, Industriepark Zwijnaarde 7, Box 4, B-9052 Ghent (BE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:
29 July 2004

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: DETECTION, IDENTIFICATION AND DIFFERENTIATION OF EUBACTERIAL TAXA USING A HYBRIDIZATION ASSAY

(57) Abstract: The present invention relates to a method for the specific detection and/or identification of *Staphylococcus* species, in particular *Staphylococcus aureus*, using new nucleic acid sequences derived from the ITS (Internal Transcribed Spacer) region. The present invention relates also to said new nucleic acid sequences derived from the ITS region, between the 16S and 23S ribosomal ribonucleic acid (rRNA) or rRNA genes, to be used for the specific detection and/or identification of *Staphylococcus* species, in particular of *S. aureus*, in a biological sample. It relates also to nucleic acid primers to be used for the amplification of said spacer region of *Staphylococcus* species in a sample.

WO 2004/052606 A3